



Worksheet-2

(C. Organic Chemistry) Hydrocarbons

- Q.1 The reaction of chlorine with methane is carried out in the presence of diffused sunlight. What is function of the light?
 - A) To break up the C H bonds in methane
 - B) To break up the chlorine molecules into free radicals
 - C) To heat up the mixture
 - D) To break up the chlorine molecules into ions
- Q.2 Methane when burnt in the presence of metallic catalyst (Cu), at high temperature (400°C) and pressure (200atm), which of the following is the ultimate product?
 - A) Methanol
- C) Methanal

B) Ethanal

- D) Methanoic acid
- Q.3 Chlorination of methane is believed to proceed through free radical mechanism. Which of the following is propagation step?
 - A) Cl-Cl \xrightarrow{hv} 2Cl
 - B) $CH_3 + Cl_2 \longrightarrow Cl + CH_3 Cl$
 - C) $CH_3 + HC1 \longrightarrow H' + H_3C-C1$
 - D) $H_3C'+C'H_3 \longrightarrow H_3C-CH_3$
- Q.4 Nitrobenzene maybe prepared by reacting benzene with a mixture of conc. H₂SO₄ and conc. HNO₃ at 55°C. Which of the following best explains the role of conc. H₂SO₄?
 - A) It removes water
 - B) It forms an unstable complex with benzene
 - C) It is protonating nitric acid
 - D) It acts as a solvent
- Q.5 Benzene reacts with acetyl chloride in the presence of catalyst AlCl₃ to give:
 - A) Aldehyde
- C) Benzyl Chloride

USE THIS SPACE FOR SCRATCH WORK

USE THIS SPACE FOR

- B) Acetophenone
- D) Benzophenone

SCRATCH WORK

- Q.6 β-elimination is competitive to nucleophilic substitution reaction. It has all of the following conditions for reaction as compared to nucleophilic substitution reaction EXCEPT:
 - A) It takes place in the presence of less polar solvent (like alcohol)
 - B) It takes place at high temperature
 - C) It requires strong nucleophile (base)
 - D) It takes place at low temperature
- Q.7 All of the following are dehydrating agents EXCEPTS:
 - A) SiO₂

- C) Al₂O₃
- B) Conc. H₂SO₄
- D) H₃PO₄
- Q.8 Which of the following is correct order of ease of dehydration of alcohols?
 - A) 1° alcohol > 2° alcohol > 3° alcohol
 - B) 3° alcohol > 2° alcohol > 1° alcohol
 - C) 2° alcohol > 1° alcohol > 3° alcohol
 - D) 3° alcohol > 1° alcohol > 2° alcohol
- Q.9 Which of the following tests is not used to distinguish between alkanes and alkenes?
 - A) Baeyar's test
- C) Cl₂(CCl₄)
- B) $Br_2(CCl_4)$
- D) Tollen's test
- Q.10 A hydrocarbon, which is a liquid at room temperature, decolourizes aqueous bromine. Which could be the molecular formula of the compound?
 - A) C_2H_2

C) C₇H₁₆

B) C_2H_4

- D) $C_{10}H_{20}$
- Q.11 Which of the following alkenes does not follow Markownikov's rule?
 - A) 1-Pentene
- C) 1-Hexene

- B) 1-Butene
- D) 2-Butene
- Q.12 Aromatic compounds burn with sooty flame because:
 - A) They have high percentage of hydrogen
 - B) They have a ring structure
 - C) They have high percentage of carbon

USE THIS SPACE FOR

D) They resist in reaction with air

Q.13 Alkanes are used as fuels. We burn them for many reasons. Which of the following is not its use?

- A) They are used to generate electricity in power stations
- B) They are used to heat our homes and cook our food
- C) They are used to provide electricity for electrolytic cell
- D) They are used to provide electricity for galvanic cell
- **Q.14** Consider the following reaction:

$$H_1C = CH_1 + HBr \rightarrow H_3C - CH_2Br$$

The mechanism of reaction is:

- A) Nucleophilic addition reactions
- B) Electrophilic addition reaction
- C) Free radical substitution
- D) Nucleophilic substitution reaction

Q.15 Which property of benzene may be directly attributed to the stability associated with its delocalized pielectrons?

- A) It has a low boiling point
- B) Its enthalpy change formation (ΔH_f) is positive
- C) It is susceptible to attack by nucleophilic reagent
- D) It tends to undergo electrophilic substitution rather addition reaction

Q.16 Among the followings the compound that can be most readily nitrated is:

- A) Benzoic acid
- C) Phenol

B) Benzene

D) Chlorobenzene

Q.17 When toluene is treated with chlorine in the presence of sunlight, which of the following is ultimate product?

- A) Benzyl chloride
- C) Benzoyl chloride
- B) Benzotrichloride
- D) Benzal dichloride
- Q.18 The most important addition reaction of alkenes forms the basis of the plastic industry. Addition polymerization is such process in which smaller molecules (monomers) repeatedly combine to form large molecular having greater molar mass (polymer) as shown:

$$H_2C = CH_2 \xrightarrow{400^{\circ}C}, \longrightarrow,_{100 \text{ atm.}} (,-H_2C - CH_2),-_n$$

Traces of $O_2(0.1\%)$ n = 1000

A good quality polythene is obtained when ethene is polymerized in the presence of:


A) Aluminium triethyl (C₂H₅)₃ only

USE THIS SPACE FOR




SCRATCH WORK

- B) Titatinum tetrachloride (TiCl₄) only
- C) $TiCl_4 + AlCl_3$
- D) $TiCl_4 + Al(C_2H_5)_3$
- Q.19 Benzene cannot undergo:
 - A) Substitution reaction
- C) Addition reaction
- B) Elimination reaction
- D) Oxidation reaction
- Q.20 All of the following statements are correct EXCEPT:
 - A) Introduction of R-group in the benzene ring in the presence of AlCl₃ is called alkylation
 - B) Introduction of acyl group in the benzene ring in the presence of AlCl₃ is called acylation
 - C) Benzene cannot undergo polymerization
 - D) Ozonolysis of benzene results in the formation of (COOH)₂
- Q.21 o- and p- directing groups have all of the following properties EXPECT:
 - A) They are electron-donating groups
 - B) They increase reactivity of mono- substituted benzene ring
 - C) They have all lone pair at the central atom of molecules expect alkyl group
 - D) Halogeno-substituted benzene is more reactive than benzene
- Q.22 All of the following methods explain stability of benzene EXPECT:
 - A) Resonance energy
 - B) Resonance method
 - C) Crystal field theory
 - D) Atomic orbital treatment
- Q.23 When different alkenes are treated with hot concentrated KMnO₄ solution, different products are obtained. Which of the following alkenes produces two moles of ketone?
 - A) $H_2C = CH_2$
- $C) R^1 R^2 C = CR^3 R^4$
- B) R CH = CH R
- D) $R^{1}R^{2}C = CH^{3}R^{4}$
- Q.24 On the oxidation of toluene by acidified KMnO₄, which of the following products is obtained?



Your STE



СНО

ds A Brighter ruture!

USE THIS SPACE FOR SCRATCH WORK B) D)

- Q.25 Which of the following methods is used to prepare ethyne on the industrial scale by?
 - A) Dehydrohalogenation of vic-dihalides
 - B) Dehalogenation of tetrahalides
 - C) Electrolysis of aqueous solution of potassium salt of unsaturated dicarboxylic acids
 - D) Reaction of calcium carbide with water
- Q.26 Kolbe's electrolytic method is used to prepare ethyne. Which of the following salts of carboxylic acid is used for this purpose?
 - A) Sodium acetate
- C) Sodium oxalate
- B) Sodium succinate
- D) Potassium maleate
- Q.27 On oxidation of ethyne with strong alkaline KMnO₄ solution, the final product formed is:
 - A) Glyoxal

C) Acetic acid

B) Glycol

- D) Oxalic acid
- Q.28 Acetaldehyde is prepared by the reaction of ethyne with water in the presence of HgSO₄/ H₂SO₄ at 75°C. Number of steps involved in this reaction is:
 - A) 1

(C) 2

B) 4

- D) 3
- Q.29 When ethyne is treated with ammonical Cu_2Cl_2 solution, then ppt of dicopper acetylide are formed. The colour of ppt is?
 - A) White

- C) Yellow
- B) Reddish brown
- D) Violet
- Q.30 When acetylene is passed under pressure over an organo-nickel catalyst at 70°C ______ is formed?
 - A) Vinyl acetylene
- C) Neoprene
- B) Di-vinyl acetylene
- D) Benzene

USE THIS SPACE FOR SCRATCH WORK

Q.31 Consider the following reaction

$$HC = CH + H_2O \xrightarrow{\text{dil. } H_2SO_4}, \longrightarrow, HgSO_4 CH_2 = CH - OH$$

$$\rightleftharpoons B \xrightarrow{V_2O_5}, \longrightarrow, [O] C$$

Which of the following is correct sequence for the product shown as B, C?

- A) CH₃CHO, CH₃COOH
- B) CH₃COCH₃, CH₃COOH
- C) CH₃CH₂OH, CH₃CHO
- D) CH₃CHO, CH₃CH₂OH

Q.32 According to atomic orbital treatment of benzene, all of the following statements are correct about benzene EXCEPT:

- A) In it each carbon atom has sp²-orbital hybridization
- B) It is cyclic hexagonal planar structure
- C) It has diffused or decolonized electron cloud
- D) It has 10 sigma bonds and 6 pi electrons

Q.33 Cyclohexane is an example of:

- A) Alicyclic hydrocarbons
- B) Aromatic hydrocarbon
- C) Aliphatic saturated hydrocarbon
- D) Aliphatic unsaturated hydrocarbon



ANSWER KEY (Worksheet-02)							
1	В	11	D	21	D	31	A
2	D	12	C	22	C	32	D
3	В	13	D	23	C	33	A
4	C	14	В	24	C		
5	В	15	D	25	D		
6	D	16	C	26	D		
7	A	17	В	27	D		
8	В	18	D	28	C		
9	D	19	В	29	В		
10	D	20	D	30	D		

ANSWERS EXPLAINED

Q.1 (B) The function of the light is to break up the chlorine molecules into free radicals such as

Cl–Cl
$$^{h\mu}$$
, \longrightarrow , Cl+ , Cl (initiation).

Halogenation is believed to proceed through **free radical substitution mechanism.** It involves the three steps such as **initiation**, **propagation and termination**.

- Q.2 (D) Ultimate product of catalytic oxidation of methane is methanoic acid.
- Q.3 (B) Propagation step is such as $CH^{\bullet,3} + Cl_2 \longrightarrow CH_3 Cl + Cl^{\bullet}$
- Q.4 (C) The role of conc. H₂SO₄ is protonating nitric acid such as

$$HONO_2 + H_2SO_4$$
 $50-55^{\circ}C$ $NO^{+,2} + HSO^{-,4} + H_2O$

Acetophenone is the organic compound with the formula C₆H₅COCH₃ (also represented by the letters **PhAc** or **BzMe**). It is the simplest aromatic ketone. This colourless,

viscous liquid is a precursor to useful resins and fragrances.

(IUPAC 1-Phenylethan-1-one) Other names:

- Methyl phenyl ketone
- Phenylethanone
- Q.6 (D) β-Elimination does not take place at low temperature, however it takes place at high temperature, in the presence of less polar solvent and in the presence of strong nucleophile.
- Q.7 (A) SiO₂ is **not dehydrating** agent while others B, C and D are used as **dehydrating agent**.
- Q.8 (B) Order of reactivity of alcohols for dehydration is such as 3° alcohol > 2° alcohol > 1° alcohol. Because the order of stability of their carbocations is as 3° carbocation > 2° carbocation > 1° carbocation.
- Q.9 (D) Tollen's test is not used to distinguish between alkanes and alkenes. Tollen's test is used to distinguish between alkenes and alkynes while alkynes having acidic hydrogen terminal alkynes give this test. By passing acetylene in the ammonical silver nitrate white ppt. of disilver acetylide are obtained as shown in the reaction.

$$HC \equiv CH+2AgNO_3+2NH_4OH \longrightarrow$$

 $AgC \equiv CAg+2NH_4NO_3+2H_2O$
white ppt

Q.10 (D) Decene (C₁₀H₂₀) is an alkene with the formula C₁₀H₂₀. It is in the liquid state. It decolourizes reddish brown aqueous bromine solution because of the presence of double bond (unsaturation). Decene contains a chain of ten carbon atoms with one double bond. There are many

isomers of decene depending on the position and geometry of the double bond.

- Q.11 (D) 2-Butene is a symmetrical molecule and does not follow Markownikov's rule. Its structure is shown as follow CH₃-CH=CH-CH₃ (symmetrical molecule).
- Q.12 (C) They have **high percentage** of carbon.
- Q.13 (D) It is **not** used to provide electricity for **galvanic cell** because **galvanic cell** is itself the source of electricity.
- Q.14 (B) Electrophilic addition reaction is shown as

$$H_2C = CH_2 + {}^{\delta +}H - Br^{\delta -} \longrightarrow H_3C - CH_2Br$$

Nucleophile Electrophile

Q.15 (D) Due to the extra resonance stability of the benzene ring, it does not undergo addition reaction in which the benzene ring resonance would be destroyed. Benzene ring by undergoing preferably electrophilic substitution retains the aromatic system. Other A, B, and C options do not fulfill the condition.

$$E - Y \longrightarrow E^{+} + Y$$

Q.16 (C) Phenol is the more reactive because OH- group is activating group, order of reactivity is as follow:

Phenol > Benzene > Chrlorobenzene > Benzoic acid. Thus phenol can be the most readily nitrated.

Q.17 (B) As a result of chlorination of benzene in the presence of sunlight

mixture of products are obtained such as benzyl chloride, benzal dichloride and benzotrichloride. Benzotrichloride is the ultimate product.

- Q.18 (D) For the better quality of polyethylene, mixture of $TiCl_4$ + $Al(C_2H_5)_3$ are used as a catalyst. It is known as catalysts.
- Q.19 (B) Benzene does not give elimination and polymerization reaction.
- Q.20 (D) Benzene reacts with ozone and gives glyoxal through benzene triozonide but not oxalic acid.
- Q.21 (D) Halogeno-substituted benzene is less reactive than benzene because its inductive effect and resonance effect are in opposite direction. Its resonance effect is slightly greater than inductive effect, because of this reason halogen group is orthopara-directing group but with more deactivation of benzene ring.
- Q.22 (C) Crystal field theory doesn't explain stability of benzene. It explain color formation by the complexes of transition metal ions .It involves d-d transition.
- Q.23 (C) When $R^1R^2C = CR^3R^4$ is treated with concentrated with KMnO₄ solution two moles of ketones are obtained.
 - $R^1R^2C = CR^3R^4 + [O] \rightarrow R^1R^2C = O$

$$+ R^3 R^4 C = O$$

- Oxidation under harsh conditions using a hot, concentrated solution of KMnO₄. Three reactions take place and different products are obtained.
- If a carbon atom is bonded to two hydrogen atoms we get oxidation to a CO₂ molecule

$$H_2C = CH_2 + 6[O] \longrightarrow CO_2 + CO_2 + 2H_2O$$

If a carbon atom is bonded to one hydrogen atom and one alkyl group we get oxidation to a
 -COOH (carboxylic acid) group
 RCH = CHR + 2[O] → RCHO +
 RCHO → RCOOH + RCOOH

- Q.24 (C) Alkyl benzenes are readily oxidized by acidified KMnO₄ or K₂Cr₂O₇. In these reactions, the alkyl groups are oxidized keeping the benzene ring intact.
 - Whatever the length of an alkyl group may be, it gives only one carboxyl group. Moreover, the colour of KMnO₄ is discharged. Therefore this reactions is used as a test for alkylbenzenes.
- Q.25 (D) In the reaction between calcium carbide and water, acetylene gas is produced on the industrial scale:

$$CaC_{2(s)} + 2H_2O_{(l)} \longrightarrow Ca(OH)_{2(s)} + C_2H_{2(g)}$$

Q.26 (D) On the electrolysis of aqueous solution of potassium maleate results in the preparation of ethyne as shown in the reaction.

HC - COOK,
$$\parallel$$
,HC - COOK + 2H₂O $\xrightarrow{\text{electrolysis}}$
,H- C \equiv C \rightarrow H + 2CO₂

 $,2KOH + H_2$

Q.27 (D) Ethyne on oxidation by strong alkaline KMnO₄ gives glyoxal followed by its further oxidation results in the formation of oxalic acid as shown in the reaction.

$$HC \equiv CH$$
, $+$ 4[O],
Ethyne, Oxidising,
(Acetylene) agent
dil. alk. $KMnO_4$,
Fehler!

Q.28 (C) When acetylene is treated with water in the presence of HgSO₄/H₂SO₄ at 75°C, No of steps involved in this reaction is 2 as shown in the reactions:

i.
$$HC \equiv CH + H^{\delta+} - OH^{\delta-}$$

 $HgSO_4$
 $H_2C = CH - O - H$
ii. $H_2C = CH - O - H \rightleftharpoons CH_3 - CH_$

Q.29 (B) When ethyne is treated with ammonical Cu₂Cl₂ solution, then ppt of dicopper acetylide are formed. The colour of ppt is reddish brown as shown in the reaction.

$$CH \equiv CH + Cu_{2}Cl_{2} + 2NH_{4}OH \rightarrow$$

$$CuC \equiv CCu + 2NH_{4}Cl + 2H_{2}O$$

$$Dicopperacetylide$$

$$(Reddish brown ppt)$$

Q.30 (D) When acetylene is passed under pressure over an organo-nickel catalyst at 70°C benzene is formed as a result of addition polymerization.

$$3HC = CH \xrightarrow{Organo-Nickel \ catalyst} \xrightarrow{70^{\circ}C}$$
Acetylene
Benzene

Q.31 (A) The correct sequence for the product is as B (CH₃CHO), C (CH₃COOH) first of all product B (ethanal) is formed which on

further oxidation gives ethanoic acid as shown in the reaction

$$\begin{split} HC &\equiv CH + H_2O \overset{dil.\ H_2SO_4}{,} \xrightarrow{,} \overset{,}{HgSO_4} \\ CH_2 &= CH - OH \longrightarrow CH_3CHO, \\ V_2O_5, \xrightarrow{,} \overset{,}{(C)} & (C) \end{split}$$

- Q.32 (D) In fact, benzene has 12 sigma bonds and 6 pi electrons.
- **Q.33** (A) Cyclohexane is an example of alicyclic hydrocarbon.

